

Multipole Expansions

The potential $\phi(\mathbf{r})$ due to a distribution of charge density $\rho(\mathbf{r}')$ within a region V is

$$\phi(\mathbf{r}) = \int_V \frac{\rho(\mathbf{r}') d^3 r'}{4\pi\epsilon_0 |\mathbf{r} - \mathbf{r}'|}$$

which we wish to approximate by a series. There are several possible useful expansions but since

$$\frac{1}{|\mathbf{r} - \mathbf{r}'|} = \frac{1}{r} \left(1 + \left(\frac{r'}{r} \right)^2 - \frac{2(\mathbf{r} \cdot \mathbf{r}')}{r^2} \right)^{-1/2}$$

when $r > r'$ a simple binomial series will converge:

$$\begin{aligned} \frac{1}{|\mathbf{r} - \mathbf{r}'|} &= \frac{1}{r} \left[1 - \frac{1}{2} \left(\frac{r'^2}{r^2} - \frac{2(\mathbf{r} \cdot \mathbf{r}')}{r^2} \right) + \frac{3}{2} \frac{1}{2} \frac{1}{2!} \left(\frac{r'^2}{r^2} - \frac{2(\mathbf{r} \cdot \mathbf{r}')}{r^2} \right)^2 + \dots \right] \\ &= \frac{1}{r} + \frac{\mathbf{r} \cdot \mathbf{r}'}{r^3} - \frac{1}{2} \frac{r'^2}{r^3} + \frac{3}{8r} \cdot \frac{4(\mathbf{r} \cdot \mathbf{r}')^2}{r^4} + \dots \\ &= \frac{1}{r} + \frac{\mathbf{r} \cdot \mathbf{r}'}{r^3} + \frac{1}{2r^5} \left[3(\mathbf{r} \cdot \mathbf{r}')^2 - (\mathbf{r} \cdot \mathbf{r})(\mathbf{r}' \cdot \mathbf{r}') \right] + \dots \\ &= \frac{1}{r} + \frac{\mathbf{r} \cdot \mathbf{r}'}{r^3} + \frac{1}{2r^5} \sum_{i,j=1}^3 x_i x_j (3x'_i x'_j - \delta_{ij} r'^2) + \dots \end{aligned}$$

where x_i ($i=1,2,3$) are the Cartesian components of \mathbf{r} and δ_{ij} is the Dirac delta function. Therefore

$$\phi(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \left(\frac{q}{r} + \frac{\mathbf{r} \cdot \mathbf{p}}{r^3} + \frac{1}{2} \frac{\sum_{ij} x_i x_j Q_{ij}}{r^5} + \dots \right)$$

where q is the total charge of the source, \mathbf{p} is the dipole moment of the distribution

$$q = \int_V \rho(\mathbf{r}') d^3 r' \quad \text{and} \quad \mathbf{p} = \int_V \mathbf{r}' \rho(\mathbf{r}') d^3 r'$$

and the nine elements of \mathbf{Q} , the Cartesian quadrupole moment tensor, are defined by

$$Q_{ij} = \int_V \rho(\mathbf{r}') (3x'_i x'_j - \delta_{ij} r'^2) d^3 r'.$$

It is easy to show that

$$\text{Tr}(\mathbf{Q}) = Q_{11} + Q_{22} + Q_{33} = 0.$$

Some authors, when the distribution has azimuthal symmetry about the z -axis and therefore

$$Q_{33} = -2Q_{11} = -2Q_{22},$$

refer to Q_{33} as “*the* quadrupole moment” which can be a little confusing.

Comments

- (i) Note that the expansion is valid only when \mathbf{r} is outside V .
- (ii) For large r usually only the first non-zero term is significant.
- (iii) The expansion can equally well be made in terms of spherical harmonics.
- (iv) The field of a distribution can be reproduced by placing suitably sized multipoles at the origin.
- (v) In general the size of the dipole and higher-order terms depend on the position of the origin.